

MULTI-ELEMENTAL ANALYSIS OF CRUDE OILS FROM NIGER DELTA NIGERIA USING INSTRUMENTAL NEUTRON ACTIVATION ANALYSIS

AGAJA S A¹, ORJI E², ALEGBEMI B³, AJISAFE J. I⁴ & BWALA S. A⁵

^{1,4}Industrial Safety and Environmental Technology, Nigeria

²Department of Physics, Federal University of Technology, Owerri, Imo State, Nigeria

³Petroleum Analysis Laboratory, Nigeria

⁵Department of General Studies, Petroleum Training Institute, Effurun, Delta State, Nigeria

ABSTRACT

Trace elemental analysis using Instrumental Neutron Activation Analysis (INAA) of Crude Oils from seven oil fields in Niger Delta, Nigeria was studied, using Ghana Research Reactor (GHARR-1) operating at 15kW at a thermal flux of $5 \times 10^{11} \text{ ncm}^{-2}\text{s}^{-1}$. From the results nine trace elements were identified: Al, Br, Ca, Cl, K, Mg, Mn, Na, and V. This study and previous ones have shown that crude oils from Nigeria are lower in metal content compare to other crude oils elsewhere around the globe. The concentration ranges of the trace elements varied significantly ($P < 0.05$) from field to field.

KEYWORDS: INAA, Crude Oils, Nigeria

INTRODUCTION

The origin of trace element in petroleum is highly correlated with the origin of petroleum itself. The origin of petroleum is complex to interpret than that of most rocks. Two different models have been put forward to explain the origin of petroleum. They are: biogenic and abiogenic models respectively (Tissot and Welte 1984, Lewan 1997). Although the biogenic organic model have been widely accepted by the petroleum industry.

The presence of trace elements like; V and Fe in crude oils, bitumen, Coal and shales was first reported by Triebs (1935). The study of trace metals in petroleum have been widely reported in literature (Ball et al, 1960, Filby and Shaty 1975, Valkovic 1978, Ellrich et al., 1985). (Trace elements in Nigeria crude oils were reported by earlier workers (Ndiokwere 1983, Oluwole et al., 1993, Adeyemo et al., 2004, Akinlua 2007 and Muhammad et al., 2003, Kpeglo et al 2015). The analysis of trace element in crude oils and its knowledge provides useful information such as: Characterization of solids and water associated with crude oils, study of heavy oils, understanding fluid interaction in sedimentary basin. Also during refining process: Control of gasoline and deposits in refineries, and process plant, control and quantification of contaminants. Many analytical techniques have been used to determine trace elements in crude oils including, Atomic Absorption spectrometry (Duyck et al; 2007), inductively-coupled plasma Atomic Emission spectroscopy (Al-swaiden, 1996) x-ray florescence (Alvarez et al 1990), neutron activation analysis (Oluwole et al 1993, Adeyemo et al, 2004, Appending et al, 2012) liquid chromatography (Khuhawar et al, 1998) and high performance liquid chromatography (Muhammed et al; 2003) INAA was used as analytical technique to study the trace element distribution in the crude oil samples because of its numerous advantage over other methods which include : very low detection limit for 30-40 elements

(parts per million or parts per billion), significant matrix independence, possibility to overcome interference in complex gamma-ray spectra (radiochemical separation) and inherent capability for high level of accuracy compared to other trace elemental analysis method (IAEA, 1987).

This work aimed at using nuclear analytical technique (INAA) to determine the concentration of trace metals in some crude oils from Niger Delta, Nigeria and to compare their element concentration with those reported in literatures by earlier workers in Nigeria and elsewhere around the world.

MATERIALS AND METHODS

Samples Collection Sites

Seven crude oil samples were collected from onshore well heads within the Niger Delta basin lat. 5.233732 and long. 6.25091 In Nigeria. The map, stratigraphy and sediment logy of Niger Delta basin was reported by Reijers (2011).

Sample Preparation

The crude oil samples were collected in clean plastic bottles (1 liter bottle). Samples were prepared by weighing 200mg on polyethylene films. They were then wrapped with the samples identifies on them. Samples were then packed into polyethylene capsule of diameter 1.60cm (rabbit capsule) and heat sealed. Standards reference material orchard leaves 1571 from the National Institute of Standards and Technology (NIST) were equally weighed as test samples.

Sample Irradiation and Counting

Samples and controls were irradiated in the Ghana Research Reactor (GHARR-1) at the Ghana Atomic Energy Commission, operating at 15kW at a thermal flux of $5 \times 10^{11} \text{ n.cm}^{-2}.\text{s}^{-1}$. Samples were transferred into irradiation sites via pneumatic transfer system at a pressure of 0.60Mpa. The irradiation was categorized according to the half-life of the element of interest as shown in the irradiation scheme in table1.

Table 1: Irradiation Schemes Short Lived Radionuclide's

Element	Target Isotope	Reaction	Product Nuclide	Half Life($t^{1/2}$)	Gamma Ray Energies (KeV)	Irradiation Time (S)	Counting Time (S)
Al	^{27}Al	$^{27}\text{Al} (n, \gamma)$	^{28}Al	2.24min	1778.9		
Ca	^{48}Ca	$^{48}\text{Ca} (n, \gamma)$	^{49}Ca	8.7min	3084.4		
Cl	^{37}Cl	$^{37}\text{Cl} (n, \gamma)$	^{38}Cl	37.3min	1642.4, 2167.5	120	600
Mg	^{26}Mg	$^{26}\text{Mg} (n, \gamma)$	^{27}Mg	9.46min	843.8, 1014.4		
Mn	^{55}Mn	$^{55}\text{Mn} (n, \gamma)$	^{56}Mn	2.58h	846.7, 1810.7 2112		
V	^{51}V	$^{51}\text{V} (n, \gamma)$	^{52}V	3.76min	1431.1		
Medium Lived Radionuclide's							
Br	^{81}Br	$^{81}\text{Br} (n, \gamma)$	^{82}Br	35.3h	554.3 776.5		
K	^{41}K	$^{41}\text{K} (n, \gamma)$	^{42}K	12.36h	1524.7	300	600
Na	^{23}Na	$^{23}\text{Na} (n, \gamma)$	^{24}Na	15.02h	1368.6 2654.1		

After the irradiation radioactivity measurement of induced radionuclide was performed by a PC-based gamma-ray spectrometry set-up. It consists of an n-type HPGe detector coupled to a computer based multi-channel analyzer (MCA) via electronic modules. The relative efficiency of detector is 25% and its energy resolution of 1.8keV at gamma-ray energy

of 1332 keV of ^{60}Co . Through appropriate choice of cooling-time, detector's dead time was controlled to be less than 10% identification of gamma-ray product radionuclide was identified through the energies and quantitative analysis of the concentration was by converting the counts as area under the photo peak by the comparator method (Landsberger 1991). Both analyses were done using the gamma-ray spectrum analysis software, ORTC MEASTRO-32. For windows model A65-B32 version 6.05 UMCBI kernel version 6.06 connections-32 version 6.06 Validation of the technique for the experimental set up was done by irradiating a standard reference material (Orchard leaves 1571) for the period of time as the sample in the same location within the reactor. The analysis of the standard reference material in table 2 shows good agreement of measured values with the certified ones. The standard specifications for Bonny light crude oil was shown in table 3.

Determination of Elements Concentration in the Samples

The calculation of trace elements concentrations in the samples was carried out by the comparator method using the same geometry, equal weights of both sample and standard with the same irradiation, decay and counting times, the concentration of the elements in the samples was determined by the expression below. (Ehmann and Diane 1991)

$$C_{\text{sample}} = C_{\text{std}} \left[\frac{A_{\text{sample}}}{A_{\text{std}}} \right]$$

Where:

C_{sample} = Unknown concentration of the element in the sample

C_{std} = Known concentration of the element in the standard

A_{sample} = Activity of the sample

A_{std} = Activity of the standard

**Table 2: Irradiation of Standard Reference Material 1571
Orchard Leaves Used for Validation**

Element	No. of Radiation	Reported Values Mgkg^{-1}	This Work Mgkg^{-1}
Al	3	Not Reported	12
Br	3	10	12
Ca	3	2.09	2.11
Cl	3	700	698
K	3	1.447wt%	1.40wt%
Mg	3	0.62wt%	0.64wt%
Mn	3	91	91
Na	3	82	81
V	3	Not Reported	37

Table 3: Specification for Nigerian Bonny Light Crude Oil

S/N		
1	Specific gravity @ 60° /60°F	0.8387 – 0.8498
2	API @ 15.55°C	35.0 – 37.0 max
3	Density @ 15.55°C	0.85 g/ml max
4	Pour point	< 40°F/4.44°C
5	Sulphur content % Wt	0.14 max

6	Color	Dark brown
7	Kinematic viscosity @ 40°	3.5 cst
8	Acid number	0.39 max
9	Reid vapour pressure (RVP)	6.52psi max
10	Water and sediment	11% max
11	Iron wt, ppm	1.00 max
12	Nickel wt, ppm	4.00 max
13	Vanadium wt, ppm	2.00 max

Expectation Derivatives From One Barrels	
Products	Gallons Per Barrels
Gasoline	19.5
Distillate fuel oil	9.2
Kerosene- type (jet fuel)	4.1
Residual fuel oil	2.3
Liquefied refined gases	1.9
Still gas	1.9
Coke	1.8
Asphalt and road oil	1.3
Petrochemical feed stocks	1.2
Lubricants	0.5
Kerosene	0.2
Others	0.3

(Source: Petroleum Analysis Laboratory, PTI Effurun)

Table 4: Trace Elements Concentrations ($\mu\text{g g}^{-1}$) in Some Crude Oil Samples from Niger Delta, Nigeria

Sample (ID)	Al	Br	Ca	Cl	K	Mg	Mn	Na	V
W1	5.9 ± 0.9	9.0 ± 1.3	775.0 ± 16.3	1431.0 ± 214.6	14.2 ± 2.1	53.0 ± 8.0	9.3 ± 1.4	675.2 ± 101.3	2.7 ± 0.4
W2	15.0 ± 2.2	9.4 ± 1.4	222.0 ± 33.2	165.4 ± 24.8	236.8 ± 35.5	143.0 ± 21.5	15.0 ± 2.2	604.9 ± 90.7	0.20 ± 0.03
W3	9.3 ± 1.4	4.0 ± 0.6	654.7 ± 98.2	209.6 ± 31.4	341.5 ± 51.2	167.0 ± 25.1	24.5 ± 3.7	146.0 ± 22.0	1.3 ± 0.2
W4	68.9 ± 10.3	5.0 ± 0.7	123.6 ± 18.5	277.1 ± 41.6	4321.0 ± 648.2	117.0 ± 17.6	12.9 ± 1.9	138.3 ± 20.8	1.5 ± 0.2
W5	45.3 ± 6.8	6.0 ± 0.9	90.8 ± 13.6	494.3 ± 0.9	2543.2 ± 381.5	88.0 ± 13.0	11.0 ± 1.7	223.9 ± 33.6	2.2 ± 0.3
W6	66.0 ± 10.0	54.4 ± 8.1	669.0 ± 10.3	797 ± 119.6	8.6 ± 1.3	85.0 ± 12.8	8.7 ± 1.3	86.0 ± 13.0	0.8 ± 0.1
W7	84.0 ± 12.6	107.5 ± 6.5	209.9 ± 31.5	7.3 ± 1.1	134.0 ± 20.0	134.0 ± 20.0	14.7 ± 2.2	96.8 ± 14.5	1.1 ± 0.2

Note: Data behind ± are standard deviations.

Table 5: Pearson Correlation for the Trace Elements

	AL	Br	Ca	Cl	K	Mg	Mn	Na	V
AL	1								
Br	0.571	1							
Ca	-0.805	-0.445	1						
Cl	-0.307	0.163	0.487	1					
K	0.326	-0.446	-0.373	-0.293	1				
Mg	-0.023	-0.210	-0.068	-0.878	-0.043	1			
Mn	-0.328	-0.404	0.350	-0.613	-0.122	0.881	1		
Na	-0.728	-0.407	0.525	0.477	-0.288	-0.319	-0.214	1	
V	-0.203	-0.384	0.462	0.645	0.252	-0.648	-0.293	0.198	1

**Table 6: Comparison of Trace Elements Reported in this
Work with Other Workers around the Globe**

Trace Elements	USA ^a	Libya ^a	Venezuela ^a	Canada ^a	Middle East ^b	China ^c	Ghana ^d	Nigeria Present Work
Al	0.30	-	1.0	-	7.0	46.4	2.96	5.9-84.0
Br	0.29	1.33	-	0.491	-	19.6	1.67	4.0-107.5
Ca	1.6	-	-	-	-	-	0.31	90.8-775.0
Cl	1.47	1.81	-	39.3	-	1000	12.95	7.3-1431.0
K	2.5	4.93	2.1	-	0.7	-	0.09	8.6-4321
Mg	2.0	-	1.7	-	7.0	-	1.32	53.0-143.0
Mn	1.2	0.79	0.21	0.048	-	5.89	0.28	8.7-24.5
Na	13.2	19.0	20.3	3.62	0.5	891	110	86.0-675.2
V	7.5	8.2	11.10	13.6	100	10.4	0.40	0.2-2.7

^a Filby and Shan 1975 and Hitchon et al 1975

^b Williams and Cawley 1963

^c Chifang et al 1991

^d Kpeglo et al 2015

**Table 7: Comparison of Vanadium Concentrations in Nigerian Crude
Oils Reported by this Work and Other Workers**

Element	Concentration	Unit	Source
V	11.2-29.2	μgg^{-1}	Akinlua et al., 2007
	0.009-1.832	μgg^{-1}	Nwachukwu et al., 1995
	3.70- 40.0	μgg^{-1}	Udo et al., 1992
	0.54-1.195	μgg^{-1}	Oluwole et al., 1993
	0.642-0.10	μgg^{-1}	Ndiokwere, 1983
	22.5-1060	ngg^{-1}	Muhammad et al., 2013
	14000-99000	ngg^{-1}	Olajire and Oderinde 1995
	1.95-7.94	μgg^{-1}	Kpeglo et al., 2015
	0.2-2.7	μgg^{-1}	Present Work

RESULTS AND DISCUSSIONS

INAA technique was used to analyze short –lived and medium-lived trace elements in crude oils from seven oil fields in Niger Delta Nigeria, nine trace elements: Al, Br, Ca, Cl, K, Mg, Mn, Na and V were analyzed and Shown in table 4. From the data it can be observed that the trace elemental concentration of the crude oils varied appreciably due to variation of geographical location of the oil fields. The nature of these trace elements and their relative abundance in crude oils can provide information about the origin, migration and maturation of petroleum as well as indicating the regional geochemical prospecting base (Chifang et al., 1991, Oluwole et al., 1993, Elirich et al., 1985).

Transition elements like Mn and V in the crude oil samples could be used to explain the origin of the crude oils from the decomposition of marine living organisms (Appenteng et al, 2012). Also their ability to form complex or chelate in a σ and π bonding form with some organic compounds (Herberhold 1972). The relativity higher concentration of elements like; Ca (90 - 775) μgg^{-1} , Cl (210 - 1431) μgg^{-1} , Na (86 - 675) μgg^{-1} in the crude oil samples suggest that the source of the crude oils from sedimentary deposition in the sea.

Table 5 is the pearson correlation computed for the trace elements. It was observed that some elements showed some significantly positive correlation with each other. Br – Al (0.6), Cl – Ca (0.5), Mn – Mg (0.9), Na –Ca (0.5), Na – K (0.5), V – Ca (0.5) and V – Cl (0.6). Table 6 showed the comparison of trace elements reported in this work with other workers around the globe. The trace elements content in the crude oils of Niger Delta Nigeria was higher than those from USA and Canada reported by Filby and Shan 1975, Hitchon et al 1975. But Vanadium content in Nigerian crude oils was very low compared to the values reported by other workers. In table 7 the Vanadium in Nigerian crude oils were compared with reported values by previous workers on Nigerian crude oils. Our values were in agreement with those reported by Nwachukwu et al 1975, Oluwole et al 1993 and Kpeglo et al., 2015.

CONCLUSIONS

The (INAA) result of this study has shown that the trace element content of Nigerian crude oils is lower compared to others reported around the globe. Relatively lower concentrations of the crude oils in V and Br is indication of good quality crude oils with minimal refining process and negative environmental impacts. Good positive correlation between Ca, Cl and Na suggest marine organic origin and sea salt component.

ACKNOWLEDGEMENTS

We sincerely appreciate the Department of Petroleum Engineering, Petroleum Training Institute, Effurun, Nigeria for supplying us with the crude oil samples across Niger Delta Nigeria. Our thanks to Mr Opata N.S and the Ghana Atomic Energy Commission (GAEC) for providing the facility for this analysis

REFERENCES

1. Filby RH and Shah KR (1975) Neutron Activation methods for trace elements in crude oils: In TF Yen (ed) The role of trace metals in petroleum, Ann Arbor, Michigan.
2. Hitchin B (1977) Geochemical links between oil fields and ore deposit in sedimentary rocks: In P Gerrard (ed) proceedings of the forum on oil and ore sediments. Imperial college London, England P 1-37.
3. Ehmann DW and Diane Ev (1991) Radiochemistry and nuclear methods of analysis, chemical analysis, Vol. 116. Wiley New York P253.
4. Elirich J. Hirner AV and Stark H (1985) Chemical Geology 48 (3) P 313-323.
5. Oluwole AF, Asubiojo OL, Nwachukwu JI, Ojo JO, Ogunsola OJ, Adenumo JA, Filby RH, Fitzgerald S, Grimm C (1993) J. Radio-Analytical and Nuclear Chem. 168 (1) P145-152.
6. Lewan MD (1984) Geochim Cosmochin Acta 48, P2231-2238.
7. Chifang C. Dingz, FUJ, Sheng G (1991) J. Radio anal Nucl. Chem., 15 (1) P177-184.
8. Shah KR, Filby RH, Haller WA (1970) J. Radio anal Nucl. Chem., 6, P185-192.
9. Landsberger S (1994) Delayed Instrumental neutron activation analysis, analysis by nuclear methods, Departments of Nuclear Engineering, University of Illinois, Wiley Chichester, P121-140
10. Adeyemo DJ, Umar IM, Jonah SA, Thomas SA, Agbaji EB, Akaho EHK (2004) J. Radio anal Nucl. Chem. 26 (1) P229-231.

11. Udo OT, Ekwere S, Abrakasa S (1992) J. Min. Geol. 28(2) P228-291.
12. Al-saleh FS, Al-Rasan GA (2008) Measurements of Radiation Level in petroleum products and wastes in Riyadh City Refinery. Journal of Environmental Radioactivity 99, P1026-1031.
13. Lee RE Jr and Von Lehmden DJ (1973). Trace metal pollution in the Environment, J. Air pollution Control Assoc. 23 P853.
14. Odebunmi EO and Adeniyi SA (2004), Characterization of crude oil and petroleum products for trace elements J. Chem. Soc. Nigeria 24, P118-121.
15. Akinlua A, Ajayi TR, Adeleke BB (2007) Organic and Inorganic geochemistry of northwestern Niger Delta oils, Geochemical Journal, 41, P271-281.
16. Nwachukwu JI, Oluwole AF, Asubiojo OI, Filby RH, Grimm CA and Fitzgerald S (1995) A geochemical evaluation of Niger Delta Oils, Geology of Deltas (Oti MN and Postma G (ed's) P287-300. A.A Balkema/Rotterdam/Bwokfield.
17. Olajire AA and Oderinde PA (1996) Finger printing of crude oil and their products. A review of the applicability of some analytical techniques, Nafta 47 (a) P259-267.
18. France MW (1964) Determination of Fe, Ni and V in petroleum oils, Anal Chem. 21 (12) P1543.
19. Appenteng MK, Golow AA, Carboo D, Adomako D, Hayford MS, Yamoah AKK, Saka D, Sarfo Dk (2012) Multi-element analysis of Ghanaian crude oils by instrumental neutron activation analysis J. Radio analytical Nucl. Chemistry 292, P1197-1206.
20. Kpeglo DO, Darko EO, Akaho EHK, Adu PAS, Abankwah E, Faanu A, Adotey DK, and Opata NS (2015) Trace element Characterization of crude oil, petroleum products and waste from the Tema oil Refinery in Ghana Using Instrumental Neutron Activation Analysis. Elixir Appl. Chem. 84, P33608-33615, www.elixirpublishers.com.
21. Ndiokwere CL (1983) Radio hem. Radio anal. Letters, 59, P201.
22. IAEA (1987) Technical Report Series No. 273 Handbook on Nuclear Activation Data.
23. Ball JS, Wenger WJ, Hyden HJ, Horr CA, Myers AT (1960) Chem. Eng Data 5, P553.
24. Williams and Cawley (1963) In Biological and Environmental effects of Arsenic, Fowler BA (ed) Elsevier Science Publishers.
25. Reijers HA (2011) Stratigraphy and Sediment logy of the Niger Delta, www.geologos.com.pl. 17, 3 P133-162.
26. Alvarez G, Martinez P and Zapata C (1990) Genetic variation in a modifier system affecting the expression of Bare mutant Drosophila Subobscura Heredity 64, 1 P55-66
27. Al-swaiden Hm, (1996) the determination of lead, nickel and vanadium in Saudi Arabian Crude Oil by sequential injection analysis/inductively coupled plasma mass spectrometer, <https://www.researchgate.net>.
28. Muhammad SA, Frew RD and Hayman AR (2013) Compound – specific isotope Analysis of Diesel Fuels in a Forensic investigation, In Aris et al, (eds) From sources to solution Springer Singapore.

29. Vukovic, Kuresevic V and Fles D (2013) Copolymers of beta-Alkoxy-styrenes and JV Phenylmaleimide Croatia Chemical Acta. <https://scholargoogle.com>.
30. Tissot and Welte (1984) Petroleum Formation Occurrence, Springer – Verlag, New York, P699.